

Models of antibody responses during HIV infection


Stanca Ciupe*, Thomas B Kepler**

*University of Louisiana at Lafayette

** Duke University Medical Center

January 11th 2010

HIV, estimated prevalence*, 2007



The boundaries and names shown and the designations used on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

Data Source: WHO/UNAIDS
Map Production: Public Health Information
and Geographic Information Systems (GIS)
World Health Organization



© WHO 2008. All rights reserved


- ❑ 22 million people have died of AIDS
 - ❑ 33 million people living with HIV/AIDS
 - ❑ 14,000 new infections every day
-

20 years of research

- | | |
|---|-------|
| ■ HIV-specific antibody test developed for blood screening | 1980s |
| ■ Detection of neutralizing antibodies against HIV | |
| ■ Primary HIV isolates are highly resistant to neutralization | 1990s |
| ■ Viruses rapidly escape neutralizing antibodies in vivo | |
| ■ Most V3 loop antibodies are strain-specific (few exceptions) | |
| ■ CD4-binding site antibodies are abundant but do not neutralize primary HIV-1 isolates (few exceptions) | |
| ■ Majority of envelope-specific antibodies are non-neutralizing and do not recognize native envelope trimer | |
| ■ Broadly neutralizing antibodies IgG1b12, 2F5, 2G12 identified | |
| ■ Interference with CD4 and coreceptor binding is main mechanism of neutralizing antibodies against gp120 | |
| ■ Structure–function analysis of gp120 confirms modes of antibody interaction | |
| ■ Importance of MPER in gp41 -- broadly neutralizing MAbs 4E10 and Z13 | |
| ■ Broadly neutralizing MAbs Ig1B12, 2G12, 2F5, 4E10 have characteristic structural features | |
| ■ Passive immunization studies -- vivo evidence of neutralization activity of broadly neutralizing MAbs | 2000s |
| ■ Virus escape to antibody neutralization occurs already in acute infection. | |
| ■ Selection in viral transmission and limit in superinfection | |

Goal of HIV vaccine

- ❑ To prevent infection
- ❑ To reduce viral loads and clinical disease progression after infection



Vaccine trials to date

■ AIDSVax

- ❑ Rocombinant gp120 protein as vaccine vector
- ❑ Induced strain-specific Ab but failed to induce broadly neutralizing Ab

■ STEP

- ❑ Induction of cellular mediated immune responses
- ❑ Recombinant adenovirus serotype 5
- ❑ Suppressed by pre-existing AD5-specific Ab
- ❑ Enhanced HIV infection

■ RV 144

- ❑ ALVAC-HIV recombinant canarypox + AIDSVax
 - ❑ Successful in reducing infectivity rates
 - ❑ Believed inefficient for clade C
-

What are the challenges?

Virus

- ❑ Extensive viral clade and sequence diversity
- ❑ Early establishment of latent viral reservoirs
- ❑ Viral evasion of humoral and cellular immune responses

Antibody

- Responses are strain-specific (ssAb)
 - No method exists to elicit broadly reactive neutralizing antibodies
 - ❑ Deleted during selection
 - ❑ Wrong conformation
 - ❑ Outcompeted
-

Depletion of interfering antibodies in chronic hepatitis C patients and vaccinated chimpanzees reveals broad cross-genotype neutralizing activity

P Zhanga, L Zhonga, EB Strublea, HW, A Kachkob, K Mihalikb, ML Virata-Theimera, HJ Alterc, S Feinstoneb, M Majorb, *PNAS*, 2009 (105).

Why do adaptive immune responses cross-react?

Karen J. Fairlie-Clarke, David M. Shuker and Andrea L. Graham
Evolutionary Applications 2009 (2).

Competition

Hypothesis


- Presence of strain-specific antibodies doesn't have to exclude the presence of broadly neutralizing ones
- Antibodies can compete with each other for
 - antigen
 - space in the lymph nodes
 - T cell conjugates and/or kinetic prolongation

Model

- Competition between the two antibody types.
 - Find the parameter region where bnAb is made inefficient by ssAb
-

Model without competition

-continuous immunization



Model without competition

-continuous immunization

Consider n strain of virus V_i , and one broadly neutralizing antibody A_0 .

$$\begin{aligned}\frac{dV_i}{dt} &= (r - K_0 A_0) V_i \\ \frac{dA_0}{dt} &= \lambda_0 \sum_{i=1}^n V_i + A_0 (a - \beta A_0)\end{aligned}$$

where $V_i(t_i) = V_0$, $A_i(t_i) = 0$, $A_0(0) = 0$.

Steady-state analysis

1. The no-infection steady state $E_0 = (0, 0, \dots, 0, 0)$ is always unstable.
2. The disease free steady state $E_1 = (0, 0, \dots, 0, a/\beta)$ is GAS when $r < K_0 \frac{a}{\beta}$.
3. The chronic steady state $E^* = (V_1^*, V_2^*, \dots, V_n^*, A_0^*)$.


$$E^* = \{(V_1^*, V_2^*, \dots, V_n^*, A_0^*) / V_i \geq 0, A_0^* > 0, V_T = A_0^* (\beta A_0^* - a), A_0^* = r/K_0\}.$$

is GAS when $r > K_0 \frac{a}{\beta}$.

Biological interpretation: When the virus replication rate is smaller than the product between the bnAb affinity rate and the antibody life span, viruses will be cleared, otherwise some persist.

Model with competition

-continuous immunization



Antibody competition model

Consider n strain of virus V_i , n strain specific antibodies A_i and one broadly neutralizing antibody A_0 .

$$\frac{dV_i}{dt} = (r - KA_i - K_0A_0)V_i$$

$$\frac{dA_i}{dt} = \lambda V_i + A_i(a - \beta A_T)$$

$$\frac{dA_0}{dt} = \lambda_0 \sum_{i=1}^n V_i + A_0(a - \beta A_T)$$

where $A_T = A_0 + \sum_{i=1}^n A_i$, $V_i(t_i) = V_0$, $A_i(t_i) = 0$, $A_0(0) = 0$.

Steady-state analysis

1. The no-infection steady state $S_0 = (0, 0, 0)$.

2. The disease free steady states

$$S_1 = \{(0, A^*, A_0^*) / A_i \geq 0, A_0 \geq 0, A_T = a/\beta\}.$$

3. Let $\emptyset \neq I \subset \{1, 2, \dots, n\}$ with $\#(I) = m \geq 1$. Then there exist steady states (V^*, A^*, A_0^*) with $A_j^*, V_j^* > 0$, for all $j \in I$ and $A_j^* = V_j^* = 0$ for all $j \notin I$ iff

$$r > \frac{m\lambda_0 K_0 + \lambda K}{m(\lambda + \lambda_0)} \frac{a}{\beta}.$$

Moreover, the non-zero components of the steady state are

$$A_j^* = \frac{\lambda}{m \lambda_0} A_0^*, A_0^* = \frac{m}{m + \frac{\lambda K}{\lambda_0 K_0}} \frac{r}{K_0},$$

$$V_j^* = \frac{\beta}{m \lambda_0} \left(\left(1 + \frac{\lambda}{\lambda_0}\right) A_0^* - \frac{a}{\beta} \right) A_0^*.$$

Observations

1. There is a unique chronic steady state when $m = n$ which is GAS when

$$r > \frac{n\lambda_0 K_0 + \lambda K}{n(\lambda + \lambda_0)} \frac{a}{\beta}.$$

2. If $r > \frac{\lambda_0 K_0 + \lambda K}{\lambda + \lambda_0} \frac{a}{\beta}$ then $r > \frac{m\lambda_0 K_0 + \lambda K}{m(\lambda + \lambda_0)} \frac{a}{\beta}$ for all $m \geq 1$.

3. If $r < \frac{n\lambda_0 K_0 + \lambda K}{n(\lambda + \lambda_0)} \frac{a}{\beta}$, then $r < \frac{m\lambda_0 K_0 + \lambda K}{m(\lambda + \lambda_0)} \frac{a}{\beta}$ for all $m \leq n$ and all virus strains will be cleared.

Biological interpretation: When the combined effect of the bnAb and ssAb exceeds the virus replication rate viruses are cleared, otherwise some persist.

Global stability results

Theorem. When $m=n$ and $r > \frac{a}{\beta} \frac{nK_0\lambda_0 + K\lambda}{n(\lambda + \lambda_0)}$ the chronic steady state is GAS.

Proof :

$$W(V_1, V_2, \dots, V_n, A_0) = \sum_{i=1}^n V_i^* \int_{V_i^*}^{V_i} \left(\frac{1}{V_i^*} - \frac{1}{\tau} \right) d\tau + \frac{1}{\lambda_0} \int_{A_0^*}^{A_0} (K_0 \tau - K_0 A_0^*) d\tau + \frac{1}{\lambda} \sum_{i=1}^n \int_{A_i^*}^{A_i} (K_0 \tau - K_0 A_i^*) d\tau$$

$gW(V_1, V_2, \dots, V_n, A_1, \dots, A_n, A_0) \geq 0$ for all $(V_1, V_2, \dots, V_n, A_1, \dots, A_n, A_0)$.

$gW(V_1, V_2, \dots, V_n, A_1, \dots, A_n, A_0) = 0$ for $(V_1, V_2, \dots, V_n, A_1, \dots, A_n, A_0) = (V_1^*, V_2^*, \dots, V_n^*, A_1^*, \dots, A_n^*, A_0^*)$.

$gW(V_1, V_2, \dots, V_n, A_1, \dots, A_n, A_0) \leq 0$ for all $(V_1, V_2, \dots, V_n, A_1, \dots, A_n, A_0)$.

$\{(V_1^*, V_2^*, \dots, V_n^*, A_1^*, \dots, A_n^*, A_0^*)\}$ is the largest invariant subset of $\{W=0\}$.

Then, from the LaSalle Invariance Principle, all solutions converge to S^* .

Global stability results

Lemma 4. Assume that $m = n$ and $r < \frac{a}{\beta} \frac{nK_0\lambda_0 + K\lambda}{n(\lambda + \lambda_0)}$. Then there are only trivial steady states ($V_i^* = 0$ for all i). Then all solutions converge to the set of steady states given by $A_T = a / \beta$.

Biological interpretation: When the combined effect of strain specific and broadly neutralizing antibody exceeds the virus replication rate, the virus is cleared, otherwise it persists.

Model prediction


For parameters

$$\frac{a}{\beta} \frac{nK_0\lambda_0 + K\lambda}{n(\lambda + \lambda_0)} < r < K_0 \frac{a}{\beta}$$


gVirus gets cleared when there is no competition between antibodies.

gVirus reaches a steady state when there is competition between SS and BN Ab.

Numerical results



Natural infection - mutations



Mutation model

Start with virus V_1 and responses A_1 and A_0 , such that

$$V_1(0) = V_{1,0} \text{ and } V_j(0) = A_j(0) = A_0(0) = A_1(0) = 0.$$

$$\frac{dV_i}{dt} = r \sum_{j=1}^n q_{ij} V_j - (KA_i + K_0 A_0) V_i$$

$$\frac{dA_i}{dt} = \lambda V_i + A_i(a - \beta A_T)$$

$$\frac{dA_0}{dt} = \lambda_0 \sum_{i=1}^n V_i + A_0(a - \beta A_T)$$

where $Q = \{q_{ij}\}_{i,j}$ is the mutation matrix: $0 \leq q_{ij} \leq 1$

$$\sum_{i=1}^n q_{ij} = 1, \text{ for all } 1 \leq j \leq n.$$

No competition

Let $V = (V_1, V_2, \dots, V_n)$ interact with A_0 .

The dynamics of the systems (1) and (2) are equivalent

$$(1) \quad \frac{dV}{dt} = (rQ - K_0 A_0 I_n) V$$

$$\frac{dA_0}{dt} = \lambda_0 V_T + A_0 (a - \beta A_0)$$

$$(2) \quad \frac{dV_T}{dt} = (r - K_0 A_0) V_T$$

$$\frac{dA_0}{dt} = \lambda_0 V_T + A_0 (a - \beta A_0)$$

when the dominant eigenvalue of $Q = \{q_{ij}\}$ is simple, with corresponding eigenvector $z \geq 0$ s.t. $Qz = z$.

For example

1. Q is irreducible. Then z is an entry wise positive vector by Perron Frobenius Theorem.


When $r < K_0 \frac{a}{\beta}$ all virus strains are cleared.

When $r > K_0 \frac{a}{\beta}$ all virus strains persists.

$$2. \quad Q = \begin{pmatrix} 1 - q_{12} & 0 & 0 & \dots & 0 & 0 \\ q_{12} & 1 - q_{23} & 0 & \dots & 0 & 0 \\ \dots & & & & & \\ 0 & 0 & 0 & q_{n-1n} & 1 \end{pmatrix}.$$

When $r < K_0 \frac{a}{\beta}$ all virus strains are cleared.

When $r > K_0 \frac{a}{\beta}$ only V_n persists.



Competition

For $n = 2$,

$$q_{11} = 1 - \alpha; q_{12} = q_{21} = \alpha; q_{22} = 1,$$

The viruses are cleared when

$$r < \Omega K_0 \frac{a}{\beta}$$

and persist when

$$r > \Omega K_0 \frac{a}{\beta},$$

$$\Omega = \frac{\frac{K}{K_0} \left(\frac{\lambda}{\lambda_0} + \frac{K_0}{K} \right) \left(\frac{\lambda}{\lambda_0} + 2 \frac{K_0}{K} \right)}{\left(1 + \frac{\lambda}{\lambda_0} \right) \left\{ \left(3 \frac{\lambda}{\lambda_0} + 4 \frac{K_0}{K} \right) - \alpha \left(2 \frac{\lambda}{\lambda_0} + 4 \frac{K_0}{K} \right) + \sqrt{\left(\left(3 \frac{\lambda}{\lambda_0} + 4 \frac{K_0}{K} \right) - \alpha \left(2 \frac{\lambda}{\lambda_0} + 4 \frac{K_0}{K} \right) \right)^2 - 8 \left(\frac{\lambda}{\lambda_0} + \frac{K_0}{K} \right) \left(\frac{\lambda}{\lambda_0} + 2 \frac{K_0}{K} \right) (1 - \alpha)^2} \right\}}$$

Model prediction



For parameters

$$\Omega K_0 \frac{a}{\beta} < r < K_0 \frac{a}{\beta}$$

gVirus gets cleared when there is no competition between antibodies.

gVirus reaches a steady state when there is competition between SS and BN Ab.

Numerical results for n strains



Conclusions and future work

- Broadly neutralizing antibodies alone can control multiple HIV infections.
 - Additional immune events directed against specific HIV viral strains weaken the immune system defense, by limiting the growth of B cells producing broadly neutralizing antibodies.
 - Under global resource limitation, HIV will be controlled only when there is no delay in a viral-specific antibody response. We know that this is not achieved in vivo.
 - Competition between HIV-strain specific and analogous antibodies might explain the immune response's ineffectiveness in controlling long term HIV infections.
-

Conclusions and future work

- Use in a vaccine trial:
 - increase the affinity of bnAB (K_0)
 - increase the antibody levels (λ_0)
 - decrease n
- Biological validation.